arXiv:hep-ph/9509287v2 21 May 1996

# Determination of CKM phases through rigid polygons of flavor SU(3) amplitudes

Amol S. Dighe [1]
Enrico Fermi Institute and Department of Physics
University of Chicago, Chicago, IL 60637

**Abstract**

Some new methods for the extraction of CKM phases $\alpha$ and $\gamma$ using flavor SU(3) symmetry are suggested. Rigid polygons are constructed in the complex plane with sides equal to the decay amplitudes of B mesons into two light (charmless) pseudoscalar mesons. These rigid polygons incorporate all the possible amplitude triangles and, being overdetermined, also serve as consistency checks and in estimating the rates of some decay modes. The same techniques also lead to numerous useful amplitude triangles when octet-singlet mixing has been taken into account and nearly physical $\eta, \eta'$ are used.

[1]Email address: asdighe@kimbark.uchicago.edu

# 1 Introduction

In the standard model, CP violation is parametrized by the Cabibbo - Kobayashi - Maskawa (CKM) [1] matrix. The unitarity of the CKM matrix implies $V_{ub}^* V_{ud} + V_{cb}^* V_{cd} + V_{tb}^* V_{td} = 0$, suggesting a unitarity triangle with its three sides as these three $V^*V$ terms and the angles $\alpha, \beta, \gamma$ ($\alpha + \beta + \gamma = \pi$) which, in Wolfenstein's parametrization [2], take the form

$$\alpha = \mathrm{Arg}(-\frac{V_{td}}{V_{ub}^*}), \quad \beta = -\mathrm{Arg}(V_{td}), \quad \gamma = \mathrm{Arg}(V_{ub}^*). \tag{1}$$

The current data from the measurements of $\frac{\epsilon'}{\epsilon}$, $B - \overline{B}$ mixing, and $\frac{|V_{ub}|}{|V_{cb}|}$ [3] give the allowed ranges of these three phases (95% c.l.) as

$$-1.00 \leq \sin 2\alpha \leq 1.00, \quad 0.21 \leq \sin 2\beta \leq 0.93, \quad 0.12 \leq \sin^2 \gamma \leq 1.00 \quad . \tag{2}$$

The decays of B mesons to light pseudoscalar mesons (B $\rightarrow$ PP) give us access to the third row and third column of the CKM matrix, where all the above phases lie. Experiments will give us the magnitudes of the amplitudes of the decay into various decay channels. (The data about the time-dependence of the decays will be available, but we shall not use that here). If theory expects the amplitudes of some three decays to form a triangle in the complex plane, constructing this triangle from the experimentally measured amplitudes will give us the relative phases between these amplitudes, from which information about the three phases above can be obtained.

Assuming flavour SU(3) symmetry gives us numerous such triangle relations and ways to determine these phases. Some such ways, with or without using any time-dependent information, have been suggested in [4]-[9] and the extent of SU(3) breaking effects has been estimated [10] to be about 20%.

The major contributions to the amplitudes of decays are from the tree or penguin type diagrams. The tree diagrams involve the process $b \rightarrow uW$ with the other quark in the B meson acting as a spectator, whereas penguins are taken to be dominated by $t$-quark exchange. (Corrections to the $t$-quark dominance of the $b \rightarrow d$ and $b \rightarrow s$ QCD penguin amplitudes [11] have been neglected in this analysis.) The phases contributed to various types of diagrams by the CKM matrix elements in the dominating term are as shown in Table (1).

| Diagram type | $\|\Delta S\| = 0$ | $\|\Delta S\| = 1$ |
|---|---|---|
| Tree | $\mathrm{Arg}(V_{ub}^* V_{ud}) = \gamma$ | $\mathrm{Arg}(V_{ub}^* V_{us}) = \gamma$ |
| Penguin | $\mathrm{Arg}(V_{tb}^* V_{td}) = -\beta$ | $\mathrm{Arg}(V_{tb}^* V_{ts}) = 0$ |

Table 1: The weak phases for tree and penguin.

Section 2 discusses the representation of decay amplitudes in terms of an SU(3) invariant basis. Section 3 gives the "rigid polygon" relations between these amplitudes. Sections 4 and 5 outline the strategies for extracting CKM phases from the relative phases of the amplitudes for $|\Delta S| = 1$ and $\Delta S = 0$ respectively. They also illustrate these methods with examples of particular decay modes and comment on their experimental feasibility. Section 6 discusses the singlet-octet $\eta$ mixing and the additional amplitudes introduced due to this. Section 7 gives some of the corresponding useful amplitude triangles when approximate physical particles $\eta, \eta'$ are used instead of $\eta_8, \eta_1$. Section 8 concludes.

# 2 Representation of amplitudes within SU(3)

Two main approaches have been taken [5, 6, 7, 12] for finding the amplitude triangle (or quadrilateral) relations. One method is to represent the amplitudes in terms of the basis of $T$ (tree), $P$ (penguin), $C$ (color-suppressed tree), $E$ (exchange), $A$ (annihilation) and $PA$ (penguin annihilation) diagram contributions. An exhaustive list of all such amplitudes has been made in [6] and some quadrangle, triangle or equivalence relations have been obained. The contributions by $E$, $A$, $PA$ have been neglected since they are expected to be suppressed by a factor of $\frac{f_B}{m_B} = 5\%$. ($E$ and $A$ will also be helicity suppressed by a factor $\frac{m_q}{m_b}$ where $q = u, d, s$).

Another equivalent approach [7, 12] is to represent the amplitudes in the basis of six SU(3) invariant amplitudes whose combinatorial coefficients will be the 6 invariant quantities formed by the combinations of the 3-vector $B_i \equiv (B^+, B^0, B_s)$, two pseudoscalar matrices $M_j^i$ (one for each pseudoscalar), and $H$, the hamiltonian for $\overline{b} \to \overline{q_1} q_2 \overline{q_3}$. $H$ can be split using SU(3) into $\overline{3} \otimes 3 \otimes \overline{3} = \overline{3} \oplus \overline{3} \oplus 6 \oplus \overline{15}$ and thus its transformation properties can be encoded into

$H^i(3)$, a vector, $H_k^{[ij]}(6)$, a traceless tensor antisymmetric in the upper two indices and $H_k^{(ij)}(15)$, a traceless tensor symmetric in the upper two indices.

The tree amplitude, in terms of this basis, will be (modulo the CKM factors)

$$\begin{aligned} T = \quad & A_3^T B_i H^i(3) M_k^j M_j^k \quad && + C_3^T B_i M_j^i M_k^j H^k(3) \\ & + A_6^T B_i H_k^{ij}(6) M_j^l M_l^k \quad && + C_6^T B_i M_j^i H_l^{jk}(6) M_k^l \\ & + A_{15}^T B_i H_k^{ij}(15) M_j^l M_l^k \quad && + C_{15}^T B_i M_j^i H_l^{jk}(15) M_k^l \quad . \end{aligned} \tag{3}$$

The $A_i^T$ are the terms that come from contracting the light quark part of the B-vector directly with the hamiltonian. This would imply that the light quark is an active part of the decay process and not just a spectator. Such amplitudes (corresponding to E, A, PA) will be suppressed by a factor of $\frac{f_B}{m_B}$ and hence can be neglected to a first approximation. The tests for this approximation (which will come naturally from the method of grouping described below) are specified at the end of this section.

The values of the nonzero elements of $H(3), H(6), H(15)$ are given in [7, 12]. The convention used for the members of the pseudoscalar meson octet is

$$\begin{aligned} \pi^+ = u\overline{d}, \qquad & \pi^0 = \tfrac{1}{\sqrt{2}}(d\overline{d} - u\overline{u}), \qquad && \pi^- = -\overline{u}d, \\ K^+ = u\overline{s}, \qquad & K^0 = d\overline{s}, \qquad && \overline{K^0} = \overline{d}s, \\ K^- = -\overline{u}s, \quad & \eta_8 = \tfrac{1}{\sqrt{6}}(2s\overline{s} - u\overline{u} - d\overline{d}). \end{aligned} \tag{4}$$

An exhaustive list of all the coefficients in $B \to PP$ is given in Table 2.

The decay modes having the same $C_i$ coefficients are grouped together and given the same group number. The unprimed (primed) group numbers correspond to the decay modes with $|\Delta S| = 0$ (1). $A(j)$ represents the amplitude for all the decay modes in that group. The advantage of using the amplitudes in terms of group numbers is that in case of multiple decay modes in a group, the ones easier to detect can be used or a suitable average of all the modes within a group can be taken to improve statistics. Comparisons of branching fractions of decay modes within a group also serve as tests of flavor SU(3) symmetry.

The *factor* column gives the factor by which the actual decay amplitude should be multiplied to get the *group* amplitude, e.g.

$$A^T(4') \equiv \sqrt{6} A^T(B^+ \to K^+ \eta_8) \quad . \tag{5}$$

| Group no. | Decay mode | | | Coefficients of $C_3$ | $C_6$ | $C_{15}$ | $A_3$ | $A_6$ | $A_{15}$ | Factor | Linear Combination |
|---|---|---|---|---|---|---|---|---|---|---|---|
| 1 | $B^+$ | $K^+$ | $\overline{K^0}$ | 1 | $-1$ | $-1$ | 0 | 1 | 3 | 1 | $p$ |
| | $B^0$ | $K^0$ | $\overline{K^0}$ | 1 | $-1$ | $-1$ | 2 | 1 | $-3$ | 1 | |
| | $B^0$ | $\pi^0$ | $\eta_8$ | 1 | $-1$ | $-1$ | 0 | 1 | $-5$ | $-\sqrt{3}$ | |
| $1'$ | $B_s$ | $K^0$ | $\overline{K^0}$ | 1 | $-1$ | $-1$ | 2 | 1 | $-3$ | 1 | $p$ |
| | $B^+$ | $K^0$ | $\pi^+$ | 1 | $-1$ | $-1$ | 0 | 1 | 3 | 1 | |
| $2'$ | $B^+$ | $K^+$ | $\pi^0$ | 1 | $-1$ | 7 | 0 | 1 | 3 | $-\sqrt{2}$ | $t+c+p$ |
| 3 | $B^+$ | $\pi^+$ | $\pi^0$ | 0 | 0 | 8 | 0 | 0 | 0 | $-\sqrt{2}$ | $t+c$ |
| $4'$ | $B^+$ | $K^+$ | $\eta_8$ | 1 | $-1$ | $-9$ | 0 | 1 | 3 | $\sqrt{6}$ | $-t-c+p$ |
| 5 | $B^+$ | $\pi^+$ | $\eta_8$ | $-2$ | 2 | $-6$ | 0 | -2 | -6 | $\sqrt{6}$ | $-t-c-2p$ |
| 6 | $B^0$ | $\pi^0$ | $\pi^0$ | 1 | 1 | $-5$ | 2 | $-1$ | 1 | $\sqrt{2}$ | $-c+p$ |
| | $B_s$ | $\overline{K^0}$ | $\pi^0$ | 1 | 1 | $-5$ | 0 | $-1$ | $-1$ | $\sqrt{2}$ | |
| | $B_s$ | $\overline{K^0}$ | $\eta_8$ | 1 | 1 | $-5$ | 0 | $-1$ | $-1$ | $\sqrt{6}$ | |
| $6'$ | $B^0$ | $K^0$ | $\pi^0$ | 1 | 1 | $-5$ | 0 | $-1$ | $-1$ | $\sqrt{2}$ | $-c+p$ |
| | $B^0$ | $K^0$ | $\eta_8$ | 1 | 1 | $-5$ | 0 | $-1$ | $-1$ | $\sqrt{6}$ | |
| $7'$ | $B_s$ | $\pi^0$ | $\eta_8$ | 0 | $-2$ | 4 | 0 | 2 | $-4$ | $-\sqrt{2}$ | $c$ |
| 8 | $B^0$ | $\pi^+$ | $\pi^-$ | 1 | 1 | 3 | 2 | $-1$ | 1 | $-1$ | $t+p$ |
| | $B_s$ | $K^-$ | $\pi^+$ | 1 | 1 | 3 | 0 | $-1$ | $-1$ | $-1$ | |
| $8'$ | $B^0$ | $K^+$ | $\pi^-$ | 1 | 1 | 3 | 0 | $-1$ | $-1$ | $-1$ | $t+p$ |
| | $B_s$ | $K^+$ | $K^-$ | 1 | 1 | 3 | 2 | $-1$ | 1 | $-1$ | |
| 9 | $B^0$ | $\eta_8$ | $\eta_8$ | 1 | $-3$ | 3 | 6 | 3 | $-3$ | $3\sqrt{2}$ | $c+p$ |
| 10 | $B^0$ | $K^+$ | $K^-$ | 0 | 0 | 0 | $-2$ | 0 | $-2$ | 1 | 0 |
| $10'$ | $B_s$ | $\pi^+$ | $\pi^-$ | 0 | 0 | 0 | 2 | 0 | 2 | $-1$ | 0 |
| | $B_s$ | $\pi^0$ | $\pi^0$ | 0 | 0 | 0 | 4 | 0 | 4 | $-1$ | |
| $11'$ | $B_s$ | $\eta_8$ | $\eta_8$ | 2 | 0 | $-6$ | 3 | 0 | $-3$ | $3/\sqrt{2}$ | $-c+2p$ |

Table 2: The coefficients of the six invariant amplitudes under SU(3).

The *linear combination* column gives the group amplitudes in terms of three complex numbers $t, c, p$; where we neglect the $A_i^T$ terms. This helps in getting the triangle relations, and will be referred to in Section 6 where the number of distinct amplitudes is greater and drawing polygon relations is not particularly instructive.

The penguin part of the amplitude can similarly be written in terms of $A_i^P$'s and $C_i^P$'s. The coefficients of $A_i^T(j), A_i^T(j'), A_i^P(j)$, and $A_i^P(j')$ have the same value and so do the coefficients of $C_i^T(j), C_i^T(j'), C_i^P(j)$, and $C_i^P(j')$. Henceforth, the superscripts $P$ and $T$ on $A_i$ and $C_i$ will be omitted wherever the relations hold true for both types of amplitudes.

The 12 amplitudes $A_i^T,\ A_i^P,\ C_i^T$, and $C_i^P$ $(i = 3, 6, 15)$ are not all independent, since there can be only 5 independent amplitudes. [When combined with the triplet light quark in the B meson, $\overline{3} \otimes 3 = 1 \oplus 8_1$, $6 \otimes 3 = 8_2 \oplus 10$, $\overline{15} \otimes 3 = 8_3 \oplus \overline{10} \oplus 27$. From PP $(8 \otimes 8 = 1 \oplus 8 \oplus 8 \oplus 10 \oplus \overline{10} \oplus 27)$, we get three symmetric states, one singlet, one (symmetrized) octet and one 27-plet. The coupling of these two implies that the decays are characterised by one singlet, three octets and one 27-plet, a total of 5 independent amplitudes.] With $A_i$ terms neglected, the number of independent amplitudes reduces to three. There are 3 relations between the six amplitudes $C_i^{T(P)}$, given by

$$C_3^T = C_6^T + C_{15}^T \quad , \quad C_6^P = C_{15}^P = 0 \quad . \tag{6}$$

The net amplitudes for the decay modes can be written as

$$A(j) \quad = A^P(j) + A^T(j) \quad = V_{tb}^* V_{td} P(j) + V_{ub}^* V_{ud} T(j) \tag{7}$$

$$A(j') \quad = A^P(j') + A^T(j') \quad = V_{tb}^* V_{ts} P(j') + V_{ub}^* V_{us} T(j') \tag{8}$$

where $P = |P|e^{i\delta_P}$ and $T = |T|e^{i\delta_T}$ are the penguin- and tree- type contributions modulo the CKM factors, and $A^P$ and $A^T$ include the CKM factors.

We clearly have the relations

$$A^P(j') = \frac{V_{ts}}{V_{td}} A^P(j) \quad , \quad A^T(j') = \frac{V_{us}}{V_{ud}} A^T(j) \tag{9}$$

between the primed and the unprimed amplitudes, but nothing can be inferred *a priori* about the relationship between $A(j)$ and $A(j')$ unless either one of $A^P$ or $A^T$ can be neglected.

Electroweak penguins do not have the same SU(3) representations as QCD penguins (the $u$ and $d$ quarks definitely interact with the photon with

different strengths). But they always appear with the $T$, $C$, and $P$ diagrams in fixed combinations (the left hand sides of equations in (10)), so the triangle relations based solely on the basis of matching of $T$, $C$, and $P$ diagrams hold true even in the presence of these electroweak penguins.

The *linear combination* column can be translated in the language of $T$, $C$, $P$, $P_{EW}$ diagrams [8] as (the superscript $C$ stands for *color-suppressed*)

$$t = T + (c_u - c_d)P^C_{EW} \ , \quad c = C + (c_u - c_d)P_{EW} \ , \quad p = P + c_d P^C_{EW} \ , \tag{10}$$

where $c_u,\ c_d,\ (c_s = c_d)$ are the strengths with which u, d, and s quarks, respectively, interact with $Z$ and $\gamma$ in electroweak penguins.

Neglecting $A^T_i, A^P_i$ contributions gives $A(10) = A(10') = 0$. This thus serves as a test for the validity of this approximation. The rates of the decays $B^0 \to K^+K^-, B_s \to \pi^+\pi^-, B_s \to \pi^0\pi^0$ should be suppressed by $\frac{f_B}{m_B}$.

# 3 The "rigid polygon" relations

An amplitude triangle is formed by three decay modes $a, b, c$ iff there exist three numbers $n_a, n_b, n_c$ such that $n_a A(a) + n_b A(b) + n_c A(c) = 0$, i.e.

$$n_a C_i(a) + n_b C_i(b) + n_c C_i(c) = 0 \quad for \quad i = 3, 6, 15 \quad , \tag{11}$$

where $C_i(a)$ is the coefficient of $C_i$ in the decay mode $a$. We shall denote a triangle formed with the sides $A(a), A(b), A(c)$ as $\triangle(a-b-c)$ where $a, b, c$ are the corresponding group numbers. All possible triangles with the $B \to PP$ decay amplitudes as their sides can be found and represented concisely in the form of two distinct rigid polygons, one each for $|\Delta S| = 1$ and $\Delta S = 0$. (See Figure 1.) The polygons are overdetermined, so the amplitudes and phases of some of the decay modes can be predicted from the others.

The polygons are oriented such that the penguin-only decay modes are along the real axis. These are the modes with $A^T = 0$ so that their net phase can be written simply as $\delta_P + \mathrm{Arg}(V^*_{tb}V_{tq})$ where $q = (d,\ s)$ for $|\Delta S| = (0,\ 1)$. When we draw these rigid polygons, we thus know the phases of all the amplitudes modulo this phase, since all the decay modes are connected via these rigid polygons to either (1) or (1′).

One major advantage of having penguin-only decay modes in the polygons is that for these decays, $|A(1)| = |\tilde{A}(1)|$ and $|A(1')| = |\tilde{A}(1')|$ where $\tilde{A}(j)$ is




Figure 1: The rigid amplitude polygons (a) $|\Delta S| = 1$ (b) $\Delta S = 0$ .

the amplitude of the CP-conjugate process in which all the particles in $(j)$ have been replaced by their antiparticles. While superposing the "particle" and "antiparticle" triangles in the process of extraction of the CKM phases, we shall always superpose their penguin-only sides.

# 4 CKM phases from $|\Delta S| = 1$ triangles

In this case, the penguin-only modes for both particle and antiparticle decay have the phase $\delta_P + \pi$, so that aligning these amplitudes along the real axis is equivalent to rotating the amplitude triangles by an unknown but fixed phase, $\delta_P$.

As a result of this rotation, the amplitude of a generic decay is now, from Eq. (8),

$$A_R(i') = -|V_{tb}^* V_{ts}||P| + |V_{ub}^* V_{us}||T|e^{i(\delta_T - \delta_P)}e^{i\gamma}, \tag{12}$$

where the subscript R stands for the amplitudes with their phase rotated such that the phase of the penguin-only amplitude is 0 or $\pi$.

The amplitude for the antiparticle decay is

$$\tilde{A}_R(i') = -|V_{tb}^* V_{ts}||P| + |V_{ub}^* V_{us}||T|e^{i(\delta_T - \delta_P)}e^{-i\gamma}. \tag{13}$$

Subtracting the two equations gets rid of the penguin contribution.

$$A_R(i') - \tilde{A}_R(i') \;\; = \;\; 2i \sin\gamma |V_{ub}^* V_{us}||T|e^{i(\delta_T - \delta_P)}$$

$$= \quad 2i \sin\gamma |A^T(i')| e^{i(\delta_T - \delta_P)} \quad . \tag{14}$$

Now, if the tree contributions to $|A^T(i')|$ and $|A^T(j)|$ can be related for some $j$ and furthermore if the penguin contribution to $A(j)$ can be neglected (as is the case whenever the tree contribution is not color-suppressed [8]), then the measurement of the decay rate gives us directly the value of $|A^T(i)|$, hence $|A^T(i')|$, and from Eq. (14) we obtain the value of $\sin\gamma$ and $\delta_T - \delta_P$.

## 4.1 $\underline{\triangle(1' - 2' - 4')}$

$$A(4') = 2A(1') - A(2') \quad . \tag{15}$$

Constructing the CP-conjugate triangle with $\tilde{A}(4'), \tilde{A}(2'), \tilde{A}(1')$ and superposing $A(1')$ and $\tilde{A}(1')$ (this is possible since $|A(1')| = |\tilde{A}(1')|$; this will introduce a discrete twofold ambiguity, leading to a discrete twofold ambiguity in $\sin\gamma$), the line joining the remaining vertices of these two triangles is

$$A_R(2') - \tilde{A}_R(2') = 2i \sin\gamma |A^T(2')| e^{i(\delta_T - \delta_P)} \quad . \tag{16}$$

Also,

$$|A^T(2')| = |A^T_{K^+\pi^0}| = \frac{|V_{us}|}{|V_{ud}|} \frac{f_K}{f_\pi} |A^T_{\pi^+\pi^0}| = \frac{|V_{us}|}{|V_{ud}|} \frac{f_K}{f_\pi} |A^T(3)| \quad . \tag{17}$$

The factor of $f_K/f_\pi$ comes from taking into account the first order SU(3) breaking under the assumption of factorization. The dominating contribution to $|A^T_{K^+\pi^0}|(|A^T_{\pi^+\pi^0}|)$ is from the tree diagram, in which $K^+(\pi^+)$ is formed purely through a weak current $[W^+ \to K^+(\pi^+)]$. With factorization, this implies a multiplicative term $< K^+|\overline{s}\gamma_\mu\gamma_5 u|0 >$ ($< \pi^+|\overline{d}\gamma_\mu\gamma_5 u|0 >$) in the amplitude, which is proportional to $f_K(f_\pi)$. This is precisely where the first order SU(3) breaking appears [10].

$B^+ \to \pi^+\pi^0$ proceeds only via $I = 2$ channel, so the QCD penguin does not contribute here. (QCD penguin is a pure $\Delta I = 1/2$ operator.) Since the electroweak penguin can be neglected in $\Delta S = 0$ channels [8], $|A^T(3)| \approx |A(3)|$ and the measurement of $|A(3)|$ along with Eq. (16) gives the value of $\sin\gamma$ and $\delta_T - \delta_P$ up to a twofold ambiguity.

If factorization holds here to a fair extent, then

$$|A_R(2') - \tilde{A}_R(2')| \leq 2|A^T(2')| \approx \frac{|V_{us}|}{|V_{ud}|} \frac{2f_K}{f_\pi} |A(3)|. \tag{18}$$

This provides a weak partial test (necessary, but not sufficient) for factorization.

The same triangle has been suggested by Deshpande and He in [7]. It is restated here for the sake of completeness and to illustrate the method. The decay modes involved here have at least one charged particle in the final states (in group $(1')$, we can choose $B^+ \to K^0\pi^+$) and the branching fractions are expected to be $\mathcal{O}(10^{-5})$, so this triangle will be experimentally easy to construct. The difficulty of separating $\eta_8$ haunts it, though (as it does all the modes in this section that involve $\eta_8$).

## 4.2 $\underline{\triangle(2' - 7' - 8')}$

We already know the phase of $A_R(2')$ from the construction of $\triangle(1' - 2' - 4')$. Now we can use

$$A(2') = A(7') + A(8') \tag{19}$$

and construct this triangle on top of the earlier one. Constructing the CP-conjugate triangle and orienting it using the information about the phase of $\tilde{A}_R(2')$ from the construction in Sec. 4.1, we get the phases of $A_R(8')$ and $\tilde{A}_R(8')$. Now

$$A_R(8') - \tilde{A}_R(8') = 2i \sin\gamma |A^T(8')| e^{i(\delta_T - \delta_P)} \tag{20}$$

and assuming factorization,

$$|A^T(8')| = |A^T_{B^0 \to K^+\pi^-}| = \frac{|V_{us}|}{|V_{ud}|}\frac{f_K}{f_\pi}|A^T_{B^0\to\pi^+\pi^-}| = \frac{|V_{us}|}{|V_{ud}|}\frac{f_K}{f_\pi}|A^T(8)| \quad . \tag{21}$$

If we can neglect the QCD penguin contribution in $A(8)$, then $|A^T(8)| \approx |A(8)|$ and we get the value of $\sin\gamma$ and $\delta_T - \delta_P$ up to a discrete twofold ambiguity. The argument about the factor $f_K/f_\pi$ in Sec. 4.1 and a similar weak partial test for factorization

$$|A_R(8') - \tilde{A}_R(8')| \leq \frac{|V_{us}|}{|V_{ud}|}\frac{2f_K}{f_\pi}|A(8)| \tag{22}$$

is valid in this case also.

Alternatively,, knowing $\gamma$ and $A_R(2'), \tilde{A}_R(2')$ from Sec. 4.1, the measurement of $|A(7')|, |\tilde{A}(7')|$ will enable us to determine $A_R(8'), \tilde{A}_R(8')$ simply by geometry.

This triangle has been suggested in [8] as a part of the quadrilateral FGOD (Fig 1) with a slightly different approach. Both the decay products in the mode ($7'$) are neutral particles. So we might come across the problem of low acceptance rate here. The branching fraction for this mode is also expected to be very small [13].

# 5 CKM phases from $\Delta S = 0$ triangles

When the polygon is oriented such that the penguin-only amplitude is along the real axis, the amplitude [Eq. (7)] of a generic decay becomes

$$A_R(j) = |V_{tb}^* V_{td}||P(j)| + |V_{ub}^* V_{ud}||T(j)|e^{i(\delta_T - \delta_P)}e^{i(\pi-\alpha)} \tag{23}$$

since $\mathrm{Arg}(\frac{V_{ub}^* V_{ud}}{V_{tb}^* V_{td}}) = \pi - \alpha$.

The corresponding antiparticle amplitude oriented in a similar manner will be

$$\tilde{A}_R(j) = |V_{tb}^* V_{td}||P(j)| + |V_{ub}^* V_{ud}||T(j)|e^{i(\delta_T - \delta_P)}e^{i(-\pi+\alpha)} \quad . \tag{24}$$

When the tree contribution in $\overline{b} \to \overline{u}u\overline{d}$ is not color suppressed, the penguin contribution is expected to be $\sim \lambda(\approx 0.2)$ times the tree contribution [8] and the $|P(j)|$ term can be neglected. The angle between these two amplitudes will then be $2\alpha$.

## 5.1 $\underline{\triangle(1-5-3)}$

$$A(3) = -A(5) - 2A(1) \tag{25}$$

and superposing the CP-conjugate triangle such that $A_R(1)$ and $\tilde{A}_R(1)$ overlap, we get (up to a discrete twofold ambiguity) the relative phases of $A_R(5)$ and $\tilde{A}_R(5)$, which is $2\alpha$.

$|A(1)|$ is expected to be very small and one might have to worry about low statistics here. But we know that $A(1) = A^P(1), A(1') = A^P(1')$ and from Eq. (9), we get

$$A(1) = \frac{V_{td}}{V_{ts}} A(1'). \tag{26}$$

If the value of $\frac{|V_{td}|}{|V_{ts}|}$ is known through other means, we can use the data from $A(1')$ to determine the magnitude of $A(1)$ to be used in the construction

of this triangle. (3) and (5) have one charged particle in their final state and a $T$ contribution (in the $\Delta S = 0$ mode) which would indicate a sizeable branching fraction and higher acceptance for both of them.

## 5.2 $\underline{\triangle(1-9-6)}$

$$A(9) = 2A(1) - A(6) \quad . \tag{27}$$

This, with its CP-conjugate triangle, gives the phases of $A_R(6)$, $\tilde{A}_R(6)$, $A_R(9)$ and $\tilde{A}_R(9)$.

This triangle relation is not directly useful for finding any of the CKM phases since the tree contribution to $A(6)$ and $A(9)$ is color-suppressed and hence of the same order of magnitude as the penguin. The penguin contribution here, therefore, cannot be neglected. But the information about the phases will be used in the construction of the next triangle.

## 5.3 $\underline{\triangle(9-5-8)}$ **or** $\underline{\triangle(6-3-8)}$

$$A(9) + A(8) = -A(5) \tag{28}$$

or

$$A(8) - A(6) = A(3), \tag{29}$$

with the information about the phase of $A_R(6)$ or $A_R(9)$ from Sec. 5.2, gives the phase of $A_R(8)$ and the CP-conjugate triangle gives the phase of $\tilde{A}_R(8)$. The phase difference between these two is $2\alpha$, similar to Sec. 5.1.

$\triangle(6-3-8)$ is the same as the $\pi - \pi$ isospin triangle in [14]. With $(3), (5), (8)$ having T contributions (and consequently, a charged particle in the final state), the branching fractions and the acceptance for these modes is expected to be on the higher side.

$\triangle(1'-6'-7')$ – a part of the quadrilateral suggested in [8] – and $\triangle(1'-6'-11')$ are possible, but are not very useful since the tree contribution to $A(7')$, $A(6')$ and $A(11')$ is color-suppressed and the electroweak penguin contributions is expected to be significant.

# 6 Physical $\eta$ and $\eta'$

The SU(3) eigenstates $\eta_8, \eta_1$ are different from the physical particles $\eta, \eta'$. Taking into account the mixing angle of $\approx 20^\circ$ [15], these physical states are very close to [6, 16]

$$\eta = \frac{1}{\sqrt{3}}(s\overline{s} - u\overline{u} - d\overline{d}) \quad , \tag{30}$$

$$\eta' = \frac{1}{\sqrt{6}}(2s\overline{s} + u\overline{u} + d\overline{d}) \quad . \tag{31}$$

Since we have contributions from the singlet component here, the trace of the pseudoscalar matrix M is no longer zero. There will, therefore, be additional terms in the amplitude, whose coefficients would have been zero had we been dealing with only the pseudoscalar octet mesons.

The additional terms in the amplitude will be

$$\begin{aligned} E_3 B_i M^i_j H^j(3) M^k_k &+ D_3 B_i H^i(3) M^j_j M^k_k \\ &+ D_6 B_i H^{ij}_k(6) M^k_j M^l_l \\ &+ D_{15} B_i H^{ij}_k(15) M^k_j M^l_l \quad . \end{aligned} \tag{32}$$

There are no terms with amplitudes $E_6$ or $E_{15}$ since their coefficients would involve $H^{ij}_j(6)$ and $H^{ij}_j(15)$ respectively, all of which are zero.

The terms $D_i$ will be suppressed by $\frac{f_B}{m_B}$ since these terms will correspond to some annihilation diagrams (for the same reason as for the $A_i$'s). So the only significant additional term we have here is the $E_3$ term.

The coefficients of $C_i$ and $E_3$ for decays involving the physical states $\eta, \eta'$ are given in Tables (3) and (4).

All the amplitudes can be explicitly written in terms of four complex numbers $t, c, p, s$ as in the *linear combination* column of Tables (2)-(3). ("$s$" is the additional amplitude due to the contribution of the singlet. It is the same as the contribution from the two-gluon diagram $P_1$ [17] and the corresponding electroweak penguin, $c_d P_{EW}$.) Thus, the knowledge of the magnitudes (obtained from the measurements) and the relative phases (obtained from the construction of triangles) of $t, c, p, s$ (or any four of their independent linear combinations) will give us the amplitudes of all the decay modes. We, therefore, need only three independent connected triangles (each triangle shares a

| Decay mode | | | Coefficients of $C_3$ | $C_6$ | $C_{15}$ | $E_3$ | Factor | Linear combination |
|---|---|---|---|---|---|---|---|---|
| $B^+$ | $K^+$ | $\eta$ | 0 | 1 | 7 | 1 | $-\sqrt{3}$ | $t+c+s$ |
| $B^+$ | $K^+$ | $\eta'$ | 3 | 1 | 1 | 4 | $\sqrt{6}$ | $t+c+3p+4s$ |
| $B_s$ | $\pi^0$ | $\eta$ | 0 | 2 | $-4$ | 0 | $\sqrt{6}$ | $-c$ |
| $B_s$ | $\pi^0$ | $\eta'$ | 0 | 2 | $-4$ | 0 | $\sqrt{3}$ | |
| $B^0$ | $K^0$ | $\eta$ | 0 | $-1$ | 3 | 1 | $-\sqrt{3}$ | $c+s$ |
| $B^0$ | $K^0$ | $\eta'$ | 3 | $-1$ | $-3$ | 4 | $\sqrt{6}$ | $c+3p+4s$ |
| $B_s$ | $\eta$ | $\eta$ | 1 | 0 | $-4$ | $-1$ | $3/\sqrt{2}$ | $-c+p-s$ |
| $B_s$ | $\eta$ | $\eta'$ | $-2$ | 0 | 5 | $-1$ | $-3/\sqrt{2}$ | $\frac{1}{2}c-2p-s$ |
| $B_s$ | $\eta'$ | $\eta'$ | 2 | 0 | $-2$ | 4 | $3/\sqrt{2}$ | $c+2p+4s$ |

Table 3: Coefficients with physical $\eta, \eta'$ for $|\Delta S| = 1$ .

| Decay mode | | | Coefficients of $C_3$ | $C_6$ | $C_{15}$ | $E_3$ | Factor | Linear combination |
|---|---|---|---|---|---|---|---|---|
| $B^+$ | $\pi^+$ | $\eta$ | 2 | $-1$ | 5 | 1 | $-\sqrt{3}$ | $t+c+2p+s$ |
| $B^+$ | $\pi^+$ | $\eta'$ | 2 | 2 | 2 | 4 | $\sqrt{6}$ | $t+c+2p+4s$ |
| $B^0$ | $\pi^0$ | $\eta$ | $-2$ | 1 | 3 | $-1$ | $\sqrt{6}$ | $-2p-s$ |
| $B^0$ | $\pi^0$ | $\eta'$ | $-1$ | $-1$ | 3 | $-2$ | $-\sqrt{3}$ | $-p-2s$ |
| $B_s$ | $\overline{K^0}$ | $\eta$ | 0 | $-1$ | 3 | 1 | $-\sqrt{3}$ | $c+s$ |
| $B_s$ | $\overline{K^0}$ | $\eta'$ | 3 | $-1$ | $-3$ | 4 | $\sqrt{6}$ | $c+3p+4s$ |
| $B^0$ | $\eta$ | $\eta$ | 1 | $-2$ | 2 | 1 | $3/\sqrt{2}$ | $c+p+s$ |
| $B^0$ | $\eta$ | $\eta'$ | 2 | $-1$ | 1 | 5 | $-3\sqrt{2}$ | $2c+2p+5s$ |
| $B^0$ | $\eta'$ | $\eta'$ | 1 | 1 | $-1$ | 4 | $3\sqrt{2}$ | $c+p+4s$ |

Table 4: Coefficients with physical $\eta, \eta'$ for $\Delta S = 0$ .

side with with at least one of the others) to get the amplitudes and phases of all the other amplitudes with the same $|\Delta S|$.

Some examples of useful triangle relations are given in Sec. 7. All the triangles have discrete twofold ambiguities associated with them.

Methods for estimating the first order SU(3) breaking effects are indicated in [10]. One way to guage the effects of SU(3) breaking on the amplitude triangles is to check if the triangle relations remain valid even when the first order SU(3) breaking terms are introduced [17]. Phase space effects have to be taken into account especially when the final state particles contain one or more heavy $\eta'$.

# 7 Amplitude triangles with $\eta$ and $\eta'$

## 7.1 $\underline{|\Delta S| = 1}$

The triangles

$$1. \quad A(B^+ \to K^0\pi^+) + \sqrt{6}A(B_s \to \pi^0\eta) = \sqrt{2}A(B^0 \to K^0\pi^0) \quad , \quad (33)$$

$$2. \quad A(B^+ \to K^0\pi^+) + \sqrt{3}A(B^0 \to K^0\eta) = \frac{3}{\sqrt{2}}A(B_s \to \eta\eta) \quad , \quad (34)$$

$$3. \quad -\sqrt{3}A(B^+ \to K^+\eta) + \tfrac{3}{\sqrt{2}}A(B_s \to \eta\eta) = -A(B^0 \to K^+\pi^-) \quad (35)$$

are three connected amplitude triangles, sufficient to predict the amplitudes and phases of all the remaining decays of the type $|\Delta S| = 1$.

The triangle in Eq. (35), when constructed on the top of the triangle in Eq. (34), gives the phase of $A_R(B^0 \to K^+\pi^-)$. The CP-conjugate $A_R(\overline{B^0} \to K^-\pi^+)$ and the method of Sec. 4.2 gives $\gamma$ and $\delta_T - \delta_P$.

All the amplitudes above have penguin contributions which are substantial in this $|\Delta S| = 1$ mode, so the branching ratios will be high, but the presence of many neutral particles in the final state might pose acceptance problems.

Some more triangles, e.g.

$$4. \quad A(B^+ \to K^0\pi^+) + \tfrac{3}{\sqrt{2}}A(B_s \to \eta'\eta') = \sqrt{6}A(B^0 \to K^0\eta') \quad , \quad (36)$$

$$5. \quad 2A(B_s \to \eta\eta) + A(B_s \to \eta'\eta') = 2A(B_s \to \eta\eta') \quad (37)$$

can be constructed which will serve to validate our assumptions collectively, if not individually. The remaining amplitudes can be generated from the information gained through Eqs. (33)-(37).

## 7.2 $\underline{\Delta S = 0}$

The triangle

$$1. \quad 3A(B^+ \to K^+\overline{K^0}) + 2\sqrt{6}A(B^0 \to \pi^0\eta) = -\sqrt{3}A(B^0 \to \pi^0\eta') \qquad (38)$$

has a penguin-only side and hence will be useful in defining the orientations of all the other amplitudes.

$$2. \quad -\sqrt{2}A(B^+ \to \pi^+\pi^0) + \sqrt{3}A(B^+ \to \pi^+\eta) = \sqrt{6}A(B^0 \to \pi^0\eta) \qquad (39)$$

constructed on top of the above triangle gives the phase of $A_R(B^+ \to \pi^+\pi^0)$. The CP-conjugate triangle gives the phase of $A_R(B^- \to \pi^-\pi^0)$ and the phase difference between these is $2\alpha$. The same procedure can be used with the information obtained from the phases of $A_R(B^+ \to \pi^+\eta)$ and $A_R(B^- \to \pi^-\eta)$, the phase difference between which is $2\alpha$.

Both $B^+ \to \pi^+\pi^0$ and $B^+ \to \pi^+\eta$ have a $T$ contribution and hence, are expected to have sizeable branching fractions and acceptances.

These triangles, along with the $\pi - \pi$ isospin triangle

$$3. \quad -\sqrt{2}A(B^+ \to \pi^+\pi^0) + \sqrt{2}A(B^0 \to \pi^0\pi^0) = -A(B^0 \to \pi^+\pi^-) \qquad (40)$$

will enable us to predict the amplitudes and phases of all the other decay modes with $\Delta S = 0$. The isospin triangle also enables one to get the phases of $A_R(B^0 \to \pi^+\pi^-)$ and $A_R(\overline{B^0} \to \pi^+\pi^-)$, the phase difference between which should be $2\alpha$.

$$4. \quad A(B^+ \to K^+\overline{K^0}) - \sqrt{3}A(B_s \to \overline{K^0}\eta) = \frac{3}{\sqrt{2}}A(B^0 \to \eta\eta) \qquad (41)$$

gives the phase of $A_R(B_s \to \overline{K^0}\eta)$, which is $\delta_T - \delta_P + \gamma$ since the only contribution here is from the $C$ diagram. (The $P_{EW}$ contribution is expected to be $\sim \lambda(\approx 0.2)$ times the $C$ contribution here [8]). The CP-conjugate triangle gives the phase of $A_R(\overline{B_s} \to K^0\eta)$, $\delta_T - \delta_P - \gamma$ and thus, $\gamma$ is obtained along with $\delta_T - \delta_P$. (This is one instance where $\delta_T - \delta_P$ is obtained

in the $\Delta S = 0$ mode. But this one will be plagued by low statistics and more neutral particles in the final state.)

Some additional triangles like

$$5.\quad 2A(B^+ \to K^+\overline{K^0}) + 3\sqrt{2}A(B^0 \to \eta'\eta') = \sqrt{6}A(B_s \to \overline{K^0}\eta') \quad , (42)$$

$$6.\quad A(B^0 \to \eta\eta) + 2A(B^0 \to \eta'\eta') = -2A(B^0 \to \eta\eta') \quad (43)$$

can be constructed for consistency checks. All the remaining amplitudes may be constructed using the information gained from the above triangles and the *linear combination* column of Table (4).

# 8 Conclusions

Using only the time-independent information about the rates of $B$ mesons decaying into light pseudoscalars, we can determine the angles of the CKM unitarity triangle under flavor SU(3) symmetry. Here we neglect the annihilation - type diagrams which are expected to be suppressed by $\frac{f_B}{m_B}$. The amplitudes are represented in terms of an SU(3) invariant basis. Rigid amplitude polygons are constructed which are overdetermined and hence can serve either for multiple ways of determining $\alpha$ and $\gamma$, as consistency checks, as tests for the approximations made, or to estimate the amplitudes for decays hard to detect experimentally. The tests for the assumptions of flavor SU(3) symmetry, factorization, annihilation diagram suppression are also built in.

The expected branching fractions of most of the decay modes are $\mathcal{O}(10^{-6}-10^{-5})$ [16] and within reach of current and upcoming experiments. The method of grouping helps in improving statistics by using the information from more than one decay mode or by allowing one to measure, say, a mode with charged decay products instead of neutral ones. The knowledge of the ratios of magnitudes of CKM elements can be used to estimate the decay rates of some modes with lower branching fractions.

The physical particles $\eta, \eta'$ are different from the SU(3) singlet $\eta_1$ or octet $\eta_8$. Taking into account this mixing, the same methods have been applied to the approximately physical $\eta, \eta'$, which will be the actual particles to be detected. The decay modes with $\eta$ or $\eta'$ as one of the decay products form a sizeable portion of charmless $B$ decays and hence taking into account the deviation of the physical states from the octet or singlet states is important. All the decay amplitudes to the approximately physical particles are expressed

explicitly in terms of four SU(3) invariant quantities and amplitude triangle relations are found which are directly useful to obtain the CKM phases, validate our assumptions and provide self-consistency tests.

# Acknowledgements

I would like to thank Aaron Grant and Mihir Worah for helpful discussions. I am particularly indebted to Jonathan Rosner for getting me interested in this topic, critically reading the manuscript and giving helpful suggestions. This work was supported in part by the U. S. Department of Energy under Contract No. DE FG02 90ER40560.

# References

[1] N. Cabibbo, Phys. Rev. Lett. **10**, 531 (1963); M. Kobayashi and T. Maskawa, Prog. Theor. Phys. **49**, 652 (1973); *B Decays*, edited by S. Stone, World Scientific, 1994.

[2] L. Wolfenstein, Phys. Rev. Lett. **13**, 562 (1964).

[3] Y. Nir and H. Quinn, Ann. Rev. Nucl. Part. Sci. **42**, 211 (1992); A. Ali and D. London, Preprint, hep-ph/9508272, 1995.

[4] D. Zeppenfeld, Z. Phys. C **8**, 77 (1981).

[5] M. Gronau, J. L. Rosner, and D. London, Phys. Rev. Lett. **73**, 21 (1994).

[6] M. Gronau, O. F. Hernandez, D. London, and J. L. Rosner, Phys. Rev. D **50**, 4529 (1994).

[7] N. G. Deshpande and X.-G. He, University of Oregon preprint OITS-576 (1995).

[8] M. Gronau, O. Hernandez, D. London, and J. L. Rosner, Preprint, Enrico Fermi Institute Report No. EFI-95-15, April, 1995, hep-ph/9504327, to be published in Phys. Rev. D.

[9] A. J. Buras and R. Fleischer, Preprint, hep-ph/9507460, 1995.

[10] M. Gronau, O. Hernandez, D. London, and J. L. Rosner, Preprint, Enrico Fermi Institute Report No. EFI-95-09, March, 1995, hep-ph/9504326, to be published in Phys. Rev. D.

[11] A. J. Buras and R. Fleischer, Phys. Lett. **B341**, 379 (1995).

[12] M. Savage and M. Wise, Phys. Rev. D **39**, 3346 (1989); **40**, 3127(E) (1989).

[13] N. G. Deshpande, X.-G. He, and J. Trampetic, Phys. Lett. **B345**, 547 (1995)

[14] M. Gronau and D. London, Phys. Rev. Lett. **65**, 3381 (1990).

[15] F. J. Gilman and R. Kauffman, Phys. Rev. D **36**, 2761 (1987); **37**, 3348(E) (1988)

[16] L. L. Chau *et al.*, Phys. Lett. D **43**, 2176 (1991).

[17] A. S. Dighe, M. Gronau, and J. L. Rosner, Phys. Lett. **B367**, 357 (1996).